

FORM B.

Return showing the description and weight of all yarn spun in the Cotton Mills in the Mysore State during the month ending September 1939

Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count
2 ^s	2,936	16½ ^s	7,537	42 ^s	30,401
4 ^s	3,961	18 ^s	4,933	44 ^s	17,190
6½ ^s	61,641	19 ^s	90,666	50 ^s	554
8 ^s	31,089	20 ^s	243,899	60 ^s	616
8½ ^s	162,669	20½ ^s	109,864	66 ^s	32
10 ^s	851	21 ^s	120,716	80 ^s	1,556
10½ ^s	152,617	22 ^s	50,285	100 ^s	12
11 ^s	37,063	22½ ^s	3,008		
11½ ^s	12,313	24 ^s	83,567		
12 ^s	33,444	26 ^s	8,425		
12½ ^s	89,963	27 ^s	7,730		
13½ ^s	13,215	28 ^s	11,270		
14 ^s	37,720	30 ^s	49,519		
14½ ^s	95,813	32 ^s	34,285		
15 ^s	20,281	36 ^s	75,157		
16 ^s	42,059	38 ^s	10,015		
		40 ^s	19,921		
				Total ...	1,778,794

BANGALORE, }
Dated 8th November 1939. }

V. VENKATACHAR,
For Director.

MYSORE STATE DEPARTMENT OF PUBLIC HEALTH.

NOTE ON THE STATE OF PUBLIC HEALTH IN MYSORE STATE
FOR THE WEEK ENDED THURSDAY, NOVEMBER 9, 1939.

A total of 31 attacks with 16 deaths from Plague was returned during the week against 32 attacks with 18 deaths in the previous week. Mysore City and Kolar Gold Field reported 5 attacks with 3 deaths, and 1 attack with 2 deaths respectively while Bangalore City continued to remain free from the disease. Comparative figures for the corresponding week in 1938 and 1937 were 58 attacks with 24 deaths and 87 attacks with 59 deaths, respectively.

A total of 1 attack with nil death from Small-pox was reported from Bangalore City.

The State remained free from Cholera, Influenza and Relapsing fever.

The total mortality for Bangalore City amounted to 117 (22 in hospitals), representing a calculated death-rate of 28·0 (22·7 excluding hospital deaths) per mille of population as against 22·3 for the corresponding week of last year. Classified according to causes, the deaths numbered—2 under Typhoid, 8 under Phthisis, 25 under Dysentery and Diarrhoea and 82 under all other causes. The reported mortality was normal.

During the week, the Civil and Military Station, Bangalore, returned 85 deaths (14 in hospitals), representing a calculated death-rate of 30·1 (25·1 excluding hospital deaths), per mille of population as against 30·1 for the corresponding week of last year. Classified according to causes, the deaths were—Broncho-Pneumonia 4, Pneumonia 4, Typhoid 1, Other Fevers 2, Phthisis 2, Dysentery and Diarrhoea 16, and all other causes 56. The reported mortality was normal.

The total mortality for Mysore City was 63 (13 in hospitals), representing a calculated death-rate of 25·9 (20·5 excluding hospital deaths) per mille of population as against 20·1 for the corresponding week of last year. Classified according to causes, the deaths were—Plague 3, Malaria 2, Other fevers 2, Phthisis 10, Dysentery and Diarrhoea 3, and all other causes 43. The reported mortality was normal.

There were 42 deaths (8 in hospitals), from all causes in Kolar Gold Field Area equivalent to a calculated death-rate of 26·3 (21·3 excluding hospital deaths) per mille of population as against 26·8 for the corresponding week of last year. Classified according to causes, the deaths numbered—2 under Plague, 2 under Malaria, 4 under other Fevers, 1 under Phthisis, 5 under Dysentery and Diarrhoea and 28 under all other causes. The reported mortality was normal.

E. R. SUNDARAJAN,
For Director.